



A generalist vector-transmitted parasite exhibits population genetic structure among host genera

Research article

Cite this article: Ellis VA, Duc M, Ciloglu A, Hellgren O, Bensch S (2025) A generalist vector-transmitted parasite exhibits population genetic structure among host genera. *Parasitology*, 1–10. <https://doi.org/10.1017/S0031182024001641>

Received: 27 June 2024
Revised: 21 November 2024
Accepted: 24 December 2024

Keywords:

avian haemosporidians; host specificity; parasite evolution; pathogen; population genomics

Corresponding author: Vincenzo A. Ellis;
Email: vaellis@udel.edu

Vincenzo A. Ellis¹ , Mélanie Duc² , Arif Ciloglu^{3,4} , Olof Hellgren⁵ and Staffan Bensch⁵

¹Department of Entomology and Wildlife Ecology, University of Delaware, Newark, DE, USA; ²P. B. Šivickis Laboratory of Parasitology, Nature Research Centre, Vilnius, Lithuania; ³Department of Parasitology, Faculty of Veterinary Medicine, Erciyes University, Kayseri, Türkiye; ⁴Vectors and Vector-Borne Diseases Implementation and Research Center, Erciyes University, Kayseri, Türkiye and ⁵Department of Biology, Lund University, Lund, Sweden

Abstract

Generalist parasites experience selective pressures from the various host species they infect. However, it is unclear if parasite transmission among host species precludes the establishment of host-specific adaptations and population genetic structure. We assessed the population genetic structure of the vector-transmitted avian haemosporidian parasite *Haemoproteus majoris* (lineage WW2; $n = 34$ infections) in a single site in southern Sweden among 10 of its host species. The 2 best-sampled host genera were *Phylloscopus* (2 species, $n = 15$ infections) and *Sylvia* (4 species, $n = 15$). We designed a sequence capture protocol to isolate 1.13 Mbp (*ca.* 5%) of the parasite genome and identified 1399 variable sites among the sequenced infections. In a principal components analysis, infections of *Phylloscopus* and *Sylvia* species mostly separated along the first 2 principal components. Sites with the highest F_{ST} values between the genera were found in genes that have mostly not been implicated in infection pathways, but several sites code for amino acid changes. An analysis of molecular variance confirmed significant variation among host genera, but not among host species within genera. The distribution of Tajima's D among sequenced loci was negatively skewed, plausibly reflecting a history of bottleneck followed by population expansion. Tajima's D was lower in infections of *Phylloscopus* than *Sylvia*, plausibly because WW2 began infecting *Phylloscopus* hosts after it was already a parasite of *Sylvia* hosts. Our results provide evidence of vector-transmitted parasite population differentiation among host species in a single location. Future work should focus on identifying the mechanisms underlying this genetic population structure.

Introduction

Host specificity is a fundamental parasite phenotype and is hypothesized to influence infectious disease emergence (Woolhouse and Gowtage-Sequeria, 2005). Parasites range from specialists that infect 1 or a few (often closely related) host species to generalists that infect many host species across varying degrees of taxonomic levels. Theoretical and empirical investigations have largely focused on testing whether host specificity is determined by evolutionary trade-offs between generalist and specialist strategies (Visher and Boots, 2020). One type of trade-off that would favour the evolution of specialists over generalists involves antagonistic pleiotropy, wherein an allele that increases a parasite's fitness in 1 host, decreases its fitness in another host (Visher and Boots, 2020). Under a scenario of antagonistic pleiotropy, one might expect a single population of a generalist to evolve into distinct genetic populations that specialize on each of the generalist's host species. The establishment of such genetic population structure might then lead to sympatric speciation of the generalist into various specialists. In the case of vector-transmitted parasites, generalists might also establish population structure because of vector biting preference. While there is evidence that generalists beget specialists over evolutionary time (Loiseau et al., 2012), closely related generalists have been identified in some groups (Mouillot et al., 2006; Ellis and Bensch, 2018) suggesting that generalists may not always evolve into specialists.

Few studies have investigated the population genetic structure of generalist parasites among their host species in the wild. In a study of a generalist nematode parasite of sympatric ungulates, Archie and Ezenwa (2011) found no population genetic differentiation in the nematode among its host species. Conversely, the seabird tick *Ixodes uriae* shows host-specific genetic population structure across its distribution (McCoy et al., 2005). Sympatric speciation and changes in host specificity, whether or not preceded by the establishment of genetic population structure, can happen rapidly. For example, the ancestor of the 2 mosquito-vectored parasites *Plasmodium*

© The Author(s), 2025. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

falciparum and *P. praefalciparum* appears to have acquired a gene through introgression (Otto *et al.*, 2018) that allowed it to infect both gorillas and humans. This was followed by a relatively small number of mutations restricting *P. falciparum* to humans (Galaway *et al.*, 2019) and its sister species (*P. praefalciparum*) to gorillas; the 2 parasites appear to have diverged about 50 kya (Otto *et al.*, 2018).

Avian haemosporidians (often called avian malaria and related parasites) are nearly globally distributed dipteran-vectored protozoan parasites (Valkiūnas, 2005). Genetic lineages of avian haemosporidians are defined by a barcoding region of the parasite cytochrome b gene (Bensch *et al.*, 2009) and exhibit great variation in host specificity (Ellis and Bensch, 2018). The avian haemosporidian species *Haemoproteus majoris* is composed of several genetic lineages including both generalists and specialists (Nilsson *et al.*, 2016). In southern Sweden, the *H. majoris* lineage WW2 frequently infects warblers from the genera *Phylloscopus* and *Sylvia* and, to a lesser degree, host species of other genera (Huang *et al.*, 2018b; Ellis *et al.*, 2020). The prevalence of WW2 is positively correlated with parasitemia among host species, and WW2 has higher prevalence and parasitemia in *Phylloscopus* and *Sylvia* warblers than in other host species (Huang *et al.*, 2018b). While avian haemosporidian vectors have often been shown to be generalist biters (e.g. Medeiros *et al.*, 2013), the vectors of WW2 in southern Sweden are unknown. However, at a similar latitude in Lithuania, the biting midges *Culicoides pictipennis*, *C. segnis* and *C. kibunensis* have been put forward as potential vectors of *H. majoris* (Žiegytė *et al.*, 2022). All 3 species have been found in southern Sweden where WW2 has been studied (unpublished data). An initial investigation of 4 parasite nuclear genes showed no genetic population structure in WW2 among its host species and little genetic variation overall (Nilsson *et al.*, 2016). However, it remains unclear if those results hold across a broader sample of the parasite genome.

Sequencing avian haemosporidian genomes is challenging because of the high abundance of host DNA in a sample of infected host blood (Videvall, 2019). Several techniques have been used to overcome this obstacle including transcriptome sequencing (Videvall *et al.*, 2017; Galen *et al.*, 2020) and genomic sequence capture (Huang *et al.*, 2018a; Barrow *et al.*, 2019; Ellis *et al.*, 2022). While transcriptome sequencing requires preserving RNA, sequence capture can be performed on previously collected DNA, making it particularly useful for retrospective analyses.

Huang *et al.* (2018b) designed sequence capture probes based on the genome of *H. tartakovskyi* and used those probes to capture and sequence part of the genome of WW2. We used the WW2 sequences from Huang *et al.* (2018b) to design new sequence capture probes targeting 1.13 Mbp of WW2 genes, *ca.* 5% of the parasite genome (Bensch *et al.*, 2016). We then performed sequence capture on WW2 infections identified in wild birds captured in southern Sweden (Huang *et al.*, 2018b; Ellis *et al.*, 2020). We used the resulting genomic data to test whether WW2 showed population genetic structure among its host species and genera and to identify the genes, if any, that show the greatest evidence of host-specific structure.

Materials and methods

Sample identification

In a previous study, we quantified the parasitemia of WW2 in avian DNA samples collected at lake Krankesjön (55°41'N, 13°26'E) in southern Sweden between 2013 and 2016 (Huang *et al.*, 2018b). We identified 48 samples with relatively high parasitemia

values and representing a diversity of host species for use in this study (host species information presented in Supplementary Table S1). Of those 48 infections, 34 passed our variant filtering (see *Bioinformatic processing* subsection) and were analysed further.

Sequence capture

Sequence capture uses biotinylated RNA probes to capture genetic sequences of interest. As a basis for probe design, we identified 1497 regions from nuclear exons (hereafter 'loci') from WW2 sequenced in a previous sequence capture study that used probes targeting *H. tartakovskyi* (GenBank BioProject accession number PRJNA448510; Huang *et al.*, 2018b). To those sequences, we added the nearly complete WW2 mitochondrial genome (5850 bp) amplified from a sample that we used for sequence capture (sample H11; Supplementary Table S1). The mitochondrial genome (GenBank accession number: PP797599) was PCR amplified and sequenced following the protocol outlined in Ciloglu *et al.* (2020). Finally, we added sequences of the caseinolytic protease C (clpC) gene of the apicoplast genome, based on the available sequences of 2 congeneric parasites, *H. magnus* (GenBank accession number: EU254647.1) and *H. passeris* (EU254653.1). We put all sequences together into a single FASTA file and provided this to TATAA Biocenter, a company that works with Arbor Biosciences to distribute myBaits probes. Arbor Biosciences removed 32 of the nuclear loci from the file, leaving 1468 targeted loci (1465 nuclear exon regions, the mitochondrial genome, and the 2 congeneric clpC sequences). The sequence names and corresponding gene identification numbers targeted from the *H. tartakovskyi* genome (Bensch *et al.*, 2016) are provided in Supplementary Table S2. We also included the latest gene annotation and descriptions in Supplementary Table S2 using the *H. tartakovskyi* annotation in version 67 of the PlasmoDB website (<https://plasmodb.org/>; Alvarez-Jarreta *et al.*, 2024); several of the genes from the original study (Bensch *et al.*, 2016) did not map or are not annotated as genes in the latest PlasmoDB version. For genes of interest with high F_{ST} values among infections in the host genera *Phylloscopus* and *Sylvia*, we provide original gene annotations and descriptions (Bensch *et al.*, 2016) if they differ from the latest annotation. We found the latest gene annotation information by mapping the originally identified genes of *H. tartakovskyi* (Bensch *et al.*, 2016) to the current PlasmoDB reference genome with BWA MEM v.0.7.17 (Li, 2013), converting the resulting SAM file to bam format with Samtools v.1.19.2 view (Danecek *et al.*, 2021) and then to bed format with bedtools v.2.30.0 bamtoBED (Quinlan and Hall, 2010) and finally extracting the gene information with bedtools intersect and further processing in R v.4.2.2 (R Core Team, 2022) with tidyverse packages (Wickham *et al.*, 2019). The FASTA file of targeted loci is provided in Supplementary File 1. Arbor Biosciences then provided us with a FASTA file of 16 850 probes, each 120 bp in length, covering our targeted sequences using 2× tiling (Supplementary File 2).

We received the synthesized probes as part of the myBaits hybridization capture for targeted NGS kit. We used this in conjunction with the Swift Biosciences Accel-NGS 2S Hyb DNA Library Kit and KAPA HiFi HotStart ReadyMix PCR Kit to perform sequence capture on our samples following the manufacturer's instructions. Briefly, we sheared the 48 DNA samples using an M220 Focused-ultrasonicator (Covaris, MA, USA) and then built indexed Illumina libraries from all samples. We next made 6 pools, each made up of 8 libraries at equimolar concentrations, and we performed the sequence capture on each pool separately. We

sequenced the captured DNA on an Illumina MiSeq (paired-end 300bp) at the Lund University DNA Sequencing Facility. All raw sequence data are available on GenBank (accession number: PRJNA1111424).

Bioinformatic processing

We assessed the quality of raw sequencing data with FastQC v.0.11.9 (<https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>). We then removed sequencing adapters and performed quality trimming using Trim Galore v.0.6.6 (<https://github.com/FelixKrueger/TrimGalore>) with the following flags: `--quality 20, --length 50`. We next mapped the paired reads to the target sequences (Supplementary File 1) using BWA MEM v.0.7.17 (Li, 2013). We used SAMtools v.1.19.2 (Danecek et al., 2021) to generate mapping statistics (SAMtools stats) and convert the resulting mapped read files to sorted bam format (SAMtools view and sort). Next, we marked duplicates in the bam files with Picard v.3.1.1 (<https://github.com/broadinstitute/picard>) MarkDuplicates and added read group information to the bam files with Picard AddOrReplaceReadGroups. We then indexed the bam files with SAMtools index and called variants with GATK v.4.3.0 (Poplin et al., 2017; Auwera and O'Connor, 2020) HaplotypeCaller with the flags `-ploidy 1` (to designate haploid samples) and `-ERC GVCF`; we prepared the reference file (Supplementary File 1) for HaplotypeCaller with SAMtools faidx and Picard CreateSequenceDictionary. We combined the resulting gzipped GVCF files with GATK CombineGVCFs followed by GATK GenotypeGVCFs. We ran most of these steps with the GNU Parallel program v.20230122 (Tang, 2023). After generating a single gzipped VCF file for all sequenced samples, we examined variant statistics in VCFtools v.0.1.16.4 (Danecek et al., 2011) and BCFtools v.1.17 (Danecek et al., 2021) and R with packages in the tidyverse. We used BCFtools to examine variants in the barcoding region of the mitochondrial genome (mtDNAgenome_CP59298, positions 4447–4925); samples with variants in that region were assumed to be mixed lineage infections and were removed from the VCF file and not analysed (all variants in that region had a minimum quality score of 30). Hard filtering VCF files is a balance between retaining samples and variable sites and removing poor quality variant calls. We filtered the VCF file with VCFtools using the following flags: `--max-missing 0.5 --minQ 30 --min-meanDP 4 --max-meanDP 250 --minDP 4 --maxDP 250`. Finally, sites in the VCF file with missing data were converted back to haploid format (from `./.` to `.`) using BCFtools fixploidy. The filtering left us with 34 samples and 1399 variable sites (1017 single nucleotide polymorphisms [SNPs] and 411 insertions/deletions [indels]; some variable sites include both SNPs and indels) among the samples in our final VCF.

Statistical analysis

We conducted a principal components analysis (PCA) in PLINK v.1.90 (Purcell et al., 2007). We started by removing variants in linkage disequilibrium calculated using 5 kb windows, a 10 bp step size, and an r^2 cut-off value of 0.20. Next, we ran the PCA on the reduced dataset and plotted the samples in multivariate space with R and packages in the tidyverse. We calculated the Weir and Cockerham weighted F_{ST} and F_{ST} per variable site among host species and genera using VCFtools (`--weir-fst-pop`). However, we downloaded a forked version of VCFtools (<https://github.com/jydu/vcftools.git>) that was modified to allow haploid

variants, which we did with the flag `-haploid`. For several variable sites with high F_{ST} between the 2 best sampled genera (*Phylloscopus* and *Sylvia*), we conducted a BLAST search of the sequenced target region (i.e. gene) on GenBank to make sure the pattern was not caused by host reads mapping to those sites. Avian genes were not among the top BLAST hits for any of the sites, and we therefore conclude that the results reflect true parasite genetic differences (results not shown). We were also interested in classifying variants as synonymous or non-synonymous. To do this we extracted consensus sequences using BCFtools consensus and organized the output into multiple sequence alignments (MSAs) using custom awk code. We then aligned the MSAs of loci with high F_{ST} between the 2 best sampled genera (*Phylloscopus* and *Sylvia*) with their corresponding genes from the *H. tartakovskyi* assembly in version 67 of the PlasmoDB website using Geneious v.11.1.5 (<http://www.geneious.com/>). Next, we extracted exon regions only from the MSAs and imported them into R as FASTA files with the R package ape v.5.7.1 (Paradis and Schliep, 2019). We then used the R package pegas v.1.3 (Paradis, 2010) to identify the haplotypes (function `haplotype()`) within each MSA and count the number of nucleotide (function `dist.dna(model = 'N')`) and amino acid (functions `trans(code = 1)` and `dist.aa()`) differences between the haplotypes (alignments were confirmed to be in the correct reading frame based using the corresponding reference gene). We calculated nucleotide diversity (π ; `--window-pi 1000`) and Tajima's D (`--TajimaD 1000`) over 1 kb windows using the forked version of VCFtools for all infections and infections of *Phylloscopus* and *Sylvia* hosts separately; we compared π and Tajima's D between infections of the 2 host genera using paired t -tests (paired because the same genomic windows were compared) with the R function `t.test()` (paired, non-parametric Wilcoxon tests gave similar results and are not reported). Finally, we read our VCF file into R with the package vcfR v.1.15.0 (Knaus and Grünwald, 2017) and conducted an analysis of molecular variance (AMOVA) in the package poppr v.2.9.6 (Kamvar et al., 2014) with the function `poppr.amova()` using the hierarchical strata of host genus and host species within genus as explanatory variables, the ade4 implementation of AMOVA, and the argument `missing = 'ignore'`. We tested the significance of the AMOVA results with the `randtest()` function with 999 randomizations (`nrepet = 999`) in the package ade4 v.1.7-22 (Chessel et al., 2004; Dray and Dufour, 2007; Dray et al., 2007; Bougeard and Dray, 2018; Thioulouse et al., 2018). We visualized the phylogenetic relationships among host species by making a consensus phylogeny of 100 trees from the posterior distribution of the analysis from `birdtree.org` (Jetz et al., 2012); we did this with the functions `consensus.edges()` in the R package phytools v.2.0-3 (Revell, 2012) and functions in ape.

Results

We identified 14 infections with variant calls in the barcoding region of the mitochondrial genome. Because such variation in that region is suggestive of mixed lineage infections (in the avian haemosporidian system, 1 nucleotide difference in the barcoding region of cytochrome b is considered a unique lineage; Bensch et al., 2009), we removed those samples from the analysis. After additional quality filtering, our final VCF file included 34 samples (i.e. infections) and 1399 variable sites (Supplementary File 3) representing infections from 10 host species and 5 host genera (Table 1; Supplementary Fig. S1). Nucleotide diversity (π) ranged from 2.85×10^{-5} to 4.04×10^{-3} (mean = $1.38 \times 10^{-4} \pm 2.21 \times 10^{-4}$ S.D.; Figure 1). Tajima's D

Table 1. Number of infections (N) of the lineage WW2, parasite *H. majoris*, analysed by host species after variant filtering and removing mixed parasite lineage infections

Host species	N
<i>Phylloscopus trochilus</i>	8
<i>Phylloscopus collybita</i>	7
<i>Sylvia communis</i>	7
<i>Sylvia atricapilla</i>	4
<i>Sylvia curruca</i>	3
<i>Sylvia borin</i>	1
<i>Acrocephalus scirpaceus</i>	1
<i>Panurus biarmicus</i>	1
<i>Parus ater</i>	1
<i>Parus palustris</i>	1

ranged from -1.84 to 2.16 (-0.29 ± 0.883 ; Figure 2). While π was higher in infections of *Phylloscopus* hosts than in infections of *Sylvia* hosts (Supplementary Fig. S2; $t = 14.03$, D.F. = 338, $P < 0.001$), Tajima's D was lower in infections of *Phylloscopus* hosts than in infections of *Sylvia* hosts (Supplementary Fig. S3; $t = -7.52$, D.F. = 292, $P < 0.001$).

The 2 best sampled host genera were *Phylloscopus* and *Sylvia* and they mostly separated in the multivariate space of a PCA along the first 2 principal component axes, representing 24.3% and 15% of the variation in the genetic data from the PCA's 20 axes, respectively (Figure 3). The lone infection of *Acrocephalus scirpaceus* also separated from the other infections (Figure 3). The loadings of the variable sites on the first 2 principal components are presented in Supplementary Table S3.


The weighted F_{ST} among all genera with more than 1 host species in the data (*Phylloscopus*, *Sylvia* and *Parus*) was 0.084 and among species with more than 1 infection (Table 1) was 0.065. The weighted F_{ST} between infections of the host genera *Phylloscopus* and *Sylvia* was 0.023. When comparing those 2 genera, there were

524 variable sites with $F_{ST} > 0$. We calculated the 95th (0.351) and 99th (0.655) percentiles of the 524 positive F_{ST} values. There were 27 F_{ST} values higher than the 95th percentile, including 6 higher than the 99th percentile (Table 2). Only 1 of these high F_{ST} values appeared to be related to infection pathways (HtGene0230, merozoite capping protein 1; Table 2). We were able to calculate the number of synonymous and non-synonymous substitutions for 6 genes (annotated as protein coding genes in the reference genome) with high F_{ST} values and found both synonymous and non-synonymous mutations (Table 3). The remaining genes could not be classified for various reasons (e.g. only 1 haplotype in the exon region of the reference, challenges with identifying the correct reading frame after alignment).

An AMOVA revealed significantly more variation than expected by chance among infections in different host genera (43.25% of the variance, $P = 0.014$) and less variation among infections of the same host species than expected by chance (52.32%, $P = 0.001$; Table 4).

Discussion

Using population genomic data from a vector transmitted generalist parasite, we found population structure among infections of different host genera, but not among infections of congeneric host species (Table 4). Infections of the 2 best sampled host genera, *Phylloscopus* and *Sylvia*, mostly separated in multivariate space of a PCA (Figure 3). The population structure was associated with genes that were largely unrelated to infection pathways (Table 2). However, 5 of 6 genes with high F_{ST} values between *Phylloscopus* and *Sylvia* hosts included non-synonymous mutations (Table 3). Lack of sufficient variation among the infections precluded more direct tests of selection (e.g. McDonald and Kreitman, 1991). Nevertheless, the large proportion of non-synonymous mutations is suggestive of selection. We therefore cannot distinguish between vector-mediated parasite isolation and host-mediated selection in causing the population genetic structure we observed. Nucleotide diversity (π ; Figure 1) and, to a lesser extent, Tajima's D (Figure 2) were negatively skewed. The latter is

**Figure 1.** Distribution of nucleotide diversity (π) in the parasite lineage WW2 calculated over 1 kb windows.




Figure 2. Distribution of Tajima's D values in the parasite lineage WW2 calculated over 1 kb windows.




Figure 3. The first 2 principal components from a PCA of genetic variation among isolates of the parasite lineage WW2 with host genus highlighted. Infections from the 2 best sampled host genera, *Phylloscopus* and *Sylvia*, mostly separate along the first 2 principal components. The variation in the genetic data explained by each principal component is presented in the axis labels. The PCA used 20 axes in total.

consistent with a demographic history of historical bottleneck in the parasite followed by population growth. We also identified several mixed lineage infections, which we did not analyse. However, future long-read sequencing may allow for analysis of such mixed lineage infections.

Parasite population structure caused by host-specific selection would plausibly be reflected in high F_{ST} values in genes involved in infection pathways, particularly genes that code for proteins that interact with host cell proteins. This may also be true of population structure among host genera since aspects of the host immune

system are evolutionary conserved (O'Connor et al., 2016). Only gene HtGene0230 had a high F_{ST} among host genera and was involved in infection pathways (Table 2). In the original annotation of the *H. tartakovskiy* reference genome (Bensch et al., 2016), this gene was annotated as merozoite capping protein 1 (MCP1). MCP1 may be involved in the attachment of the merozoite to host red blood cells (Klotz et al., 1989). However, in the latest annotation of the *H. tartakovskiy* genome, the sequence is no longer annotated as a gene (Table 2). Despite the paucity of infection pathway genes contributing to WW2's population structure among *Phylloscopus*

Table 2. Variable sites of the parasite lineage WW2 with F_{ST} values between infections in the hosts of the genera *Phylloscopus* and *Sylvia*. Only variable sites with F_{ST} values greater than the 95th percentile of all positive F_{ST} values are shown and sites with F_{ST} values greater than the 99th percentile (column 99%) are indicated as 'outliers'. Target refers to the targeted sequence that the sequence capture probes were designed to capture; POS is the position on that sequence that was variable; HtGene is the name of the gene in the original published assembly of the reference *H. tartakovskyi* genome; RefGene is the corresponding gene name in the latest assembly; type indicates whether the region is characterized as a protein coding gene or pseudogene in the latest assembly; gene function is the annotated description of the gene in the latest assembly, and the note 'original annotation': refers to the annotated description from the original assembly; both the latest and the original assembly descriptions are provided if the gene annotation differs between the assemblies. Missing RefGene and type values indicate that the gene from the original assembly is not annotated in the latest assembly. The F_{ST} values for the variable sites are also shown

Target	POS	F_{ST}	99%	HtGene	RefGene	Type	Gene function
seq73__HtGene0129	161	0.693	Outlier	HtGene0129			Original annotation: Putative uncharacterized protein OS = Plasmodium
seq75__HtGene0131	1147	0.351		HtGene0131	Htart_000018100	Pseudogene	DNA primase large subunit putative
seq93__HtGene0183	1003	0.678	Outlier	HtGene0183	Htart_000022800	Protein coding gene	Metacaspase-like protein; Original annotation: Putative uncharacterized protein OS = Plasmodium
seq114__HtGene0230	517	0.529		HtGene0230			Original annotation: Merozoite capping protein 1, putative
seq120__HtGene0243	950	0.365		HtGene0243	Htart_000029900	Protein coding gene	Methionine aminopeptidase 1b putative
seq255__HtGene0476	126	0.408		HtGene0476	Htart_000054900	Protein coding gene	TFIIH basal transcription factor complex helicase XPD subunit; Original annotation: DNA excision-repair helicase, putative
seq258__HtGene0481	71	0.382		HtGene0481	Htart_000055400	Protein coding gene	V-type proton ATPase subunit E putative
seq608__HtGene1356	382	0.360		HtGene1356	Htart_000141500	Pseudogene	Elongation factor 2
seq664__HtGene1491	78	0.680	Outlier	HtGene1491	Htart_000154400	Pseudogene	Pre-mRNA-splicing factor 38A putative; Original annotation: Putative uncharacterized protein OS = Plasmodium
seq665__HtGene1491	483	0.703	Outlier	HtGene1491	Htart_000154400	Pseudogene	Pre-mRNA-splicing factor 38A putative; Original annotation: Putative uncharacterized protein OS = Plasmodium
seq790__HtGene1778	212	0.367		HtGene1778			Original annotation: Apicoplast ribosomal protein L10, putative
seq927__HtGene2105	261	0.409		HtGene2105	Htart_000212900	Protein coding gene	LCCL domain-containing protein
seq956__HtGene2156	339	0.351		HtGene2156	Htart_000217300	Protein coding gene	Exoribonuclease putative
seq1151__HtGene2635	473	0.648		HtGene2635	Htart_000261100	Protein coding gene	Vacuolar proton translocating ATPase subunit A putative
seq1168__HtGene2682	72	0.463		HtGene2682	Htart_000266200	Pseudogene	Myosin C
seq1187__HtGene2704	433	0.462		HtGene2704	Htart_000268400	Protein coding gene	CCR4-associated factor 1; Original annotation: Putative uncharacterized protein (Fragment)
seq1234__HtGene2808	407	0.351		HtGene2808			Original annotation: RNA binding protein, putative OS = Plasmodium
seq1413__HtGene3214	1546	0.608		HtGene3214			Original annotation: Putative uncharacterized protein OS = Medicago
seq1487__HtGene3342	99	0.363		HtGene3342	Htart_000331000	Pseudogene	DNA gyrase subunit A
seq1487__HtGene3342	100	0.363		HtGene3342	Htart_000331000	Pseudogene	DNA gyrase subunit A

(Continued)

Table 2. (Continued.)

Target	POS	F_{ST}	99%	HtGene	RefGene	Type	Gene function
seq1526__HtGene3475	517	0.657	Outlier	HtGene3475	Htart_000342800	Protein coding gene	Mitosis protein dim1 putative
seq1533__HtGene3504	31	0.462		HtGene3504	Htart_000345400	Protein coding gene	40S ribosomal protein S7 putative
seq1664__HtGene3763	45	0.682	Outlier	HtGene3763	Htart_000368400	Protein coding gene	FeS cluster assembly protein SufD putative
seq1703__HtGene3892	36	0.370		HtGene3892	Htart_000381900	Protein coding gene	Zinc finger protein putative; Original annotation: Putative uncharacterized protein OS = Plasmodium
seq1703__HtGene3892	39	0.370		HtGene3892	Htart_000381900	Protein coding gene	Zinc finger protein putative; Original annotation: Putative uncharacterized protein OS = Plasmodium
seq1704__HtGene3893	1435	0.351		HtGene3893	Htart_000382000	Protein coding gene	Heat shock protein 90 putative
seq1718__HtGene3926	609	0.351		HtGene3926	Htart_000384900	Pseudogene	6-phosphofructokinase

Table 3. Target loci (target) with high F_{ST} values among *Phylloscopus* and *Sylvia* hosts (Table 2) were aligned with exons in the corresponding protein coding genes in the reference *H. tartakovskyi* assembly (RefGene); we were able to determine nucleotide substitution types (synonymous or non-synonymous) among haplotypes of 6 genes. We present the number of haplotypes per gene, the haplotype sequence length, mean and maximum number of nucleotide substitutions among haplotypes, and the corresponding mean and maximum number of amino acid substitutions in the translated haplotype sequences

Target	RefGene	Sequence length (bp)	Number of haplotypes	Nucleotide substitutions (mean/max)	Amino acid substitutions (mean/max)	Nucleotide substitution types
seq93__HtGene0183	Htart_000022800	1819	4	(1/1)	(0/0)	Synonymous
seq120__HtGene0243	Htart_000029900	1054	5	(22.9/52)	(9.4/19)	Synonymous and non-synonymous
seq927__HtGene2105	Htart_000212900	1188	3	(1.33/2)	(1.33/2)	Non-synonymous
seq1151__HtGene2635	Htart_000261100	549	2	(1/1)	(1/1)	Non-synonymous
seq1187__HtGene2704	Htart_000268400	893	5	(2/3)	(1.6/3)	Synonymous and non-synonymous
seq1664__HtGene3763	Htart_000368400	1569	3	(1.33/2)	(0.66/1)	Synonymous and non-synonymous

Table 4. Results of analysis of molecular variance (AMOVA). the sources of variation were infections among host genera ('host genera'), infections among host species within host genera ('host species within genera'), and infections in host species ('infections within host species'). Degrees of freedom (DF), sum of square deviations (SSD), mean square deviations (MSD), variance and its percent of the total variance; phi is the population differentiation statistic. The p value for each source of variance from a randomization test is also presented. Infections differed among host genera more than expected by chance and infections within host species differed less than expected by chance

	D.F.	SSD	MSD	Variance	Percent variance	Phi	P
Host genera	4	11778.69	2944.67	445.50	43.25	0.433	0.014
Host species within genera	5	3536.83	707.37	45.60	4.43	0.078	0.235
Infections within host species	24	12934.66	538.94	538.94	52.32	0.477	0.001

and *Sylvia* hosts (Table 2), 5 of 6 investigated genes included non-synonymous nucleotide substitutions separating their haplotypes (Table 3). Parasites may also experience selection in genes in response to different host physiologies, and those genes may not necessarily be related to infection pathways. Another possibility is that genes with high F_{ST} among host genera are linked to infection related genes that we did not sequence. Besides host-specific selection, vectors of WW2 may have different preferences for bird genera; we cannot distinguish between these 2 possibilities with our data. Furthermore, host selection and vector preference may be operating simultaneously. Our results emphasize the need to better study vectors in this system and to characterize more genes

in WW2's genome. Characterizing parasite gene transcription during infection (Videvall et al., 2017) and among host species (in the case of generalists) remains an important endeavour (Garcia-Longoria et al., 2020). Furthermore, we sequenced relatively high intensity infections and it is unclear if lower intensity infections would show the same patterns.

Tajima (1989) examined the relationship between the number of segregating sites and the average number of pairwise differences between sequences of individuals in a neutrally evolving population. The D statistic was described as the difference between the pairwise nucleotide differences and the number of segregating sites (the latter scaled by sample size) divided by the standard deviation

of that difference (Tajima, 1989). Negative values of D suggest more rare alleles than expected under a neutrally evolving population. A large negative D might result from a gene under selection, but here we found negative values between -1 and 0 of D across many of the genes surveyed (Figure 2). This result would appear consistent with a historical bottleneck followed by population expansion, since a growing population could produce an excess of new mutations (Hartl, 2020). While it is unclear what could have caused this, one can speculate. The closest relative to WW2 and the other *H. majoris* lineages might be a specialist of the host *Sylvia borin* (e.g. SYBOR15 was used as the closest relative of *H. majoris* in an analysis by Nilsson et al., 2016). Such a switch from specialist to generalist may have caused a bottleneck, followed by rapid population growth as the parasite began infecting additional host species. Indeed, Tajima's D values were lower in infections of *Phylloscopus* hosts than in infections of *Sylvia* hosts (Supplementary Fig. S3) as might be expected if *Phylloscopus* hosts are more recent hosts of WW2. Testing for a relationship between evolutionary shifts in host specificity and parasite population genetics awaits better supported parasite phylogenies (perhaps through mitochondrial genome sequencing; Pacheco et al., 2018; Ciloglu et al., 2020) and additional population genomics studies of related parasites. Furthermore, deeper sequencing would allow for quantification of haplotype diversity within individual infections (Videvall et al., 2017); such diversity may also be relevant for understanding the relationship between host shifts and parasite speciation.

We have demonstrated genetic population structure of a generalist parasite among the genera of its hosts. The mechanism underlying this population structure remains unknown. Our results suggest that parasites are able to diverge and perhaps eventually speciate in sympatry, although isolation among host species in sympatry is not necessarily equivalent to sympatric speciation (Pérez-Tris et al., 2007). This result is important for considering modes of parasite speciation (Pérez-Tris et al., 2007; Ricklefs et al., 2014). Future work will need to establish the relationship between parasite population structure and speciation, the frequency with which generalist parasites show signals of genetic population structure among their host species in sympatry, and the mechanisms underpinning such genetic population structure. The field of avian haemosporidian evolutionary ecology has grown rapidly over the last 2 decades with the development of DNA barcoding of parasite lineages (Bensch et al., 2000; Hellgren et al., 2004) and the establishment of a unifying database to facilitate comparative analyses (Bensch et al., 2009). Novel genomic techniques like sequence capture hold promise for again advancing the field to allow for informative tests of novel hypotheses and a better general understanding of vector-transmitted parasite evolution.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S0031182024001641>.

Data availability statement. All raw sequence data are available on GenBank (accession number: PRJNA111424).

Acknowledgements. We thank Shawn Polson (Center for Bioinformatics and Computational Biology, University of Delaware) for conducting an initial analysis of a subset of the data. We thank Tomas Johansson (Lund University) for performing the sequencing. The variant calling and filtering pipelines included elements of code from Shawn Polson's analysis and code for other pipelines written by Amelia Harrison and Madolyn MacDonald (Center for Bioinformatics and Computational Biology, University of Delaware) and from several online tutorials; VAE is very appreciative of all the examples and help. VAE also thanks Eloisa H. Reis Sari for discussing and providing insights into

the results of the study. We thank two reviewers for providing helpful feedback on an earlier version of the manuscript.

Author contributions. VAE and SB designed the study. VAE, MD, AC, OH, and SB contributed to data collection. VAE performed bioinformatic and statistical analyses and wrote the first draft of the manuscript and all authors contributed to and approved the final version of the manuscript.

Financial support. We performed bioinformatic and statistical analyses on the University of Delaware's BIOMIX cluster, which is supported by Delaware INBRE (NIH/NIGMS P20 GM103446), NIH Shared Instrumentation Grant (S10OD028725), the State of Delaware, and the Delaware Biotechnology Institute. VAE was supported by funding from USDA Hatch (DEL00774, DEL00854, and NE1943) and from the Carl Tryggers Foundation (CTS16:53). The study was also supported by the Swedish Research Council through grants to OH (VR 2016-03419 and 2021-03663) and to SB (VR 2017-03937).

Competing interests. The authors declare there are no conflicts of interest.

Ethical standards. Not applicable.

References

- Alvarez-Jarreta J, Amos B, Aurrecochea C, Bah S, Barba M, Barreto A, Basenko EY, Belnap R, Blevins A, Böhme U, Brestelli J, Brown S, Callan D, Campbell LI, Christophides GK, Crouch K, Davison HR, DeBarry JD, Demko R, Doherty R, Duan Y, Dundore W, Dyer S, Falke D, Fischer S, Gajria B, Galdi D, Giraldo-Calderón GI, Harb OS, Harper E et al. (2024) VEUPathDB: The eukaryotic pathogen, vector and host bioinformatic resource center in 2023. *Nucleic Acids Research* **52**, D808–D816. doi:10.1093/nar/gkad1003
- Archie EA and Ezenwa VO (2011) Population genetic structure and history of a generalist parasite infecting multiple sympatric host species. *International Journal for Parasitology* **41**, 89–98. doi:10.1016/j.ijpara.2010.07.014
- Auvera GAVD and O'Connor BD (2020) *Genomics in the Cloud: using Docker, GATK, and WDL in Terra*, 1st edn. USA: O'Reilly Media.
- Barrow LN, Allen JM, Huang X, Bensch S and Witt CC (2019) Genomic sequence capture of haemosporidian parasites: Methods and prospects for enhanced study of host–parasite evolution. *Molecular Ecology Resources* **19**, 400–410. doi:10.1111/1755-0998.12977
- Bensch S, Canbäck B, DeBarry JD, Johansson T, Hellgren O, Kissinger JC, Palinauskas V, Videvall E and Valkiūnas G (2016) The genome of *Haemoproteus tartakovskiy* and its relationship to human malaria parasites. *Genome Biology and Evolution* **8**, 1361–1373. doi:10.1093/gbe/evw081
- Bensch S, Hellgren O and Pérez-Tris J (2009) MalAvi: A public database of malaria parasites and related haemosporidians in avian hosts based on mitochondrial cytochrome *b* lineages. *Molecular Ecology Resources* **9**, 1353–1358. doi:10.1111/j.1755-0998.2009.02692.x
- Bensch S, Stjernman M, Hasselquist D, Örjan Ö, Hannson B, Wester Dahl H and Pinheiro RT (2000) Host specificity in avian blood parasites: A study of *Plasmodium* and *Haemoproteus* mitochondrial DNA amplified from birds. *Proceedings of the Royal Society of London. Series B: Biological Sciences* **267**, 1583–1589. doi:10.1098/rspb.2000.1181
- Bougeard S and Dray S (2018) Supervised multiblock analysis in R with the ade4 package. *Journal of Statistical Software* **86**. doi:10.18637/jss.v086.i01
- Chessel D, Dufour A and Thioulouse J (2004) The ade4 package - I: One-table methods. *R News* **4**, 5–10.
- Ciloglu A, Ellis VA, Duc M, Downing PA, Inci A and Bensch S (2020) Evolution of vector transmitted parasites by host switching revealed through sequencing of *Haemoproteus* parasite mitochondrial genomes. *Molecular Phylogenetics & Evolution* **153**, 106947. doi:10.1016/j.ympev.2020.106947
- Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G, Marth GT, Sherry ST, McVean G and Durbin R and 1000 Genomes Project Analysis Group (2011) The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158. doi:10.1093/bioinformatics/btr330

- Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane T, McCarthy SA, Davies RM and Li H (2021) Twelve years of SAMtools and BCFtools. *GigaScience* **10**, giab008. doi:10.1093/gigascience/giab008
- Dray S and Dufour A-B (2007) The ade4 package: Implementing the duality diagram for ecologists. *Journal of Statistical Software* **22**. doi:10.18637/jss.v022.i04
- Dray S, Dufour A and Chessel D (2007) The ade4 package - II: Two-table and k-table methods. *R News* **7**, 47–52.
- Ellis VA and Bensch S (2018) Host specificity of avian haemosporidian parasites is unrelated among sister lineages but shows phylogenetic signal across larger clades. *International Journal for Parasitology* **48**, 897–902. doi:10.1016/j.ijpara.2018.05.005
- Ellis VA, Huang X, Westerdahl H, Jönsson J, Hasselquist D, Neto JM, Nilsson J, Nilsson J, Hegemann A, Hellgren O and Bensch S (2020) Explaining prevalence, diversity and host specificity in a community of avian haemosporidian parasites. *Oikos* **129**, 1314–1329. doi:10.1111/oik.07280
- Ellis VA, Kalbskopf V, Ciloglu A, Duc M, Huang X, Inci A, Bensch S, Hellgren O and Palinauskas V (2022) Genomic sequence capture of *Plasmodium relictum* in experimentally infected birds. *Parasites and Vectors* **15**, 267. doi:10.1186/s13071-022-05373-w
- Galaway F, Yu R, Constantinou A, Prugnolle F and Wright GJ (2019) Resurrection of the ancestral RH5 invasion ligand provides a molecular explanation for the origin of *P. falciparum* malaria in humans. *PLOS Biology* **17**, e3000490. doi:10.1371/journal.pbio.3000490
- Galen SC, Borner J, Williamson JL, Witt CC and Perkins SL (2020) Metatranscriptomics yields new genomic resources and sensitive detection of infections for diverse blood parasites. *Molecular Ecology Resources* **20**, 14–28. doi:10.1111/1755-0998.13091
- Garcia-Longoria L, Palinauskas V, Ilgūnas M, Valkiūnas G and Hellgren O (2020) Differential gene expression of *Plasmodium homocircumflexum* (lineage pCOLL4) across two experimentally infected passerine bird species. *Genomics* **112**, 2857–2865. doi:10.1016/j.ygeno.2020.03.025
- Hartl DL (2020) *A Primer of Population Genetics and Genomics*, 4th edn. Oxford, UK: Oxford University Press.
- Hellgren O, Waldenström J and Bensch S (2004) A new PCR assay for simultaneous studies of *Leucocytozoon*, *Plasmodium*, and *Haemoproteus* from avian blood. *Journal of Parasitology* **90**, 797–802. doi:10.1645/GE-184R1
- Huang X, Ellis VA, Jönsson J and Bensch S (2018b) Generalist haemosporidian parasites are better adapted to a subset of host species in a multiple host community. *Molecular Ecology* **27**, 4336–4346. doi:10.1111/mec.14856
- Huang X, Hansson R, Palinauskas V, Valkiūnas G, Hellgren O and Bensch S (2018a) The success of sequence capture in relation to phylogenetic distance from a reference genome: A case study of avian haemosporidian parasites. *International Journal for Parasitology* **48**, 947–954.
- Jetz W, Thomas GH, Joy JB, Hartmann K and Mooers AO (2012) The global diversity of birds in space and time. *Nature* **491**, 444–448. doi:10.1038/nature11631
- Kamvar ZN, Tabima JF and Grünwald NJ (2014) Poppr: An R package for genetic analysis of populations with clonal, partially clonal, and/or sexual reproduction. *PeerJ* **2**, e281. doi:10.7717/peerj.281
- Klotz FW, Hadley TJ, Aikawa M, Leech J, Howard RJ and Miller LH (1989) A 60-kDa *Plasmodium falciparum* protein at the moving junction formed between merozoite and erythrocyte during invasion. *Molecular and Biochemical Parasitology* **36**, 177–185. doi:10.1016/0166-6851(89)90190-4
- Knaus BJ and Grünwald NJ (2017) VCFR: A package to manipulate and visualize variant call format data in R. *Molecular Ecology Resources* **17**, 44–53. doi:10.1111/1755-0998.12549
- Li H (2013) Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
- Loiseau C, Harrigan RJ, Robert A, Bowie RCK, Thomassen HA, Smith TB and Sehgal RNM (2012) Host and habitat specialization of avian malaria in Africa. *Molecular Ecology* **21**, 431–441. doi:10.1111/j.1365-294X.2011.05341.x
- McCoy KD, Chapuis E, Tirard C, Boulinier T, Michalakakis Y, Bohec CL, Maho YL and Gauthier-Clerc M (2005) Recurrent evolution of host-specialized races in a globally distributed parasite. *Proceedings of the Royal Society B: Biological Sciences* **272**, 2389–2395. doi:10.1098/rspb.2005.3230
- McDonald JH and Kreitman M (1991) Adaptive protein evolution at the *Adh* locus in *Drosophila*. *Nature* **351**, 652–654. doi:10.1038/351652a0
- Medeiros MCI, Hamer GL and Ricklefs RE (2013) Host compatibility rather than vector–host–encounter rate determines the host range of avian *Plasmodium* parasites. *Proceedings of the Royal Society B: Biological Sciences* **280**, 20122947. doi:10.1098/rspb.2012.2947
- Mouillot D, Krasnov BR, Shenbrot GI, Gaston KJ and Poulin R (2006) Conservatism of host specificity in parasites. *Ecography* **29**, 596–602. doi:10.1111/j.0906-7590.2006.04507.x
- Nilsson E, Taubert H, Hellgren O, Huang X, Palinauskas V, Markovets MY, Valkiūnas G and Bensch S (2016) Multiple cryptic species of sympatric generalists within the avian blood parasite *Haemoproteus majoris*. *Journal of Evolutionary Biology* **29**, 1812–1826. doi:10.1111/jeb.12911
- O'Connor EA, Strandh M, Hasselquist D, Nilsson J-Å and Westerdahl H (2016) The evolution of highly variable immunity genes across a passerine bird radiation. *Molecular Ecology* **25**, 977–989. doi:10.1111/mec.13530
- Otto TD, Gilabert A, Crellen T, Böhme U, Arnathau C, Sanders M, Oyola SO, Okouga AP, Boundenga L, Willaume E, Ngoubangoye B, Moukoudoum ND, Paupy C, Durand P, Rougeron V, Ollomo B, Renaud F, Newbold C, Berriman M and Prugnolle F (2018) Genomes of all known members of a *Plasmodium* subgenus reveal paths to virulent human malaria. *Nature Microbiology* **3**, 687–697. doi:10.1038/s41564-018-0162-2
- Pacheco MA, Matta NE, Valkiūnas G, Parker PG, Mello B, Stanley CE, Lentino M, Garcia-Amado MA, Cranfield M, Kosakovsky Pond SL and Escalante AA (2018) Mode and rate of evolution of haemosporidian mitochondrial genomes: Timing the radiation of avian parasites. *Molecular Biology and Evolution* **35**, 383–403. doi:10.1093/molbev/msx285
- Paradis E (2010) pegas: An R package for population genetics with an integrated–modular approach. *Bioinformatics* **26**, 419–420. doi:10.1093/bioinformatics/btp696
- Paradis E and Schliep K (2019) ape 5.0: An environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics* **35**, 526–528. doi:10.1093/bioinformatics/bty633
- Pérez-Tris J, Hellgren O, Krizanauskienė A, Waldenström J, Secondi J, Bonneaud C, Fjeldså J, Hasselquist D and Bensch S (2007) Within-host speciation of malaria parasites. *PLoS ONE* **2**, e235. doi:10.1371/journal.pone.0000235
- Poplin R, Ruano-Rubio V, DePristo MA, Fennell TJ, Carneiro MO, Van Der Auwera GA, Kling DE, Gauthier LD, Levy-Moonshine A, Roazen D, Shakir K, Thibault J, Chandran S, Whelan C, Lek M, Gabriel S, Daly MJ, Neale B, MacArthur DG and Banks E (2017) *Scaling Accurate Genetic Variant Discovery to Tens of Thousands of Samples*. bioRxiv, 201178. doi:10.1101/201178.
- Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, Maller J, Sklar P, De Bakker PIW, Daly MJ and Sham PC (2007) PLINK: A tool set for whole-genome association and population-based linkage analyses. *The American Journal of Human Genetics* **81**, 559–575. doi:10.1086/519795
- Quinlan AR and Hall IM (2010) BEDTools: A flexible suite of utilities for comparing genomic features. *Bioinformatics* **26**, 841–842. doi:10.1093/bioinformatics/btq033
- R Core Team (2022) *R: A Language and Environment for Statistical Computing*.
- Revell LJ (2012) phytools: An R package for phylogenetic comparative biology (and other things). *Methods in Ecology and Evolution* **3**, 217–223. doi:10.1111/j.2041-210X.2011.00169.x
- Ricklefs RE, Outlaw DC, Svensson-Coelho M, Medeiros MCI, Ellis VA and Latta S (2014) Species formation by host shifting in avian malaria parasites. *Proceedings of the National Academy of Sciences* **111**, 14816–14821. doi:10.1073/pnas.1416356111
- Tajima F (1989) Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics* **123**, 585–595. doi:10.1093/genetics/123.3.585
- Tang O (2023) GNU Parallel 20230122 ('Bolsonaristas'). *Zenodo*. doi:10.5281/zenodo.7558957
- Thioulouse J, Dray S, Dufour A, Siberchicot A, Jombart T and Pavoine S (2018) *Multivariate Analysis of Ecological Data with Ade4*. New York, NY: Springer.

- Valkiūnas G** (2005) *Avian Malaria Parasites and Other Haemosporidia*. USA: CRC Press.
- Videvall E** (2019) Genomic advances in avian malaria research. *Trends in Parasitology* **35**, 254–266. doi:10.1016/j.pt.2018.12.005
- Videvall E, Cornwallis CK, Ahrén D, Palinauskas V, Valkiūnas G and Hellgren O** (2017) The transcriptome of the avian malaria parasite *Plasmodium ashfordi* displays host-specific gene expression. *Molecular Ecology* **26**, 2939–2958. doi:10.1111/mec.14085
- Visher E and Boots M** (2020) The problem of mediocre generalists: Population genetics and eco-evolutionary perspectives on host breadth evolution in pathogens. *Proceedings of the Royal Society B: Biological Sciences* **287**, 20201230. doi:10.1098/rspb.2020.1230
- Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, Grolemund G, Hayes A, Henry L, Hester J, Kuhn M, Pedersen T, Miller E, Bache S, Müller K, Ooms J, Robinson D, Seidel D, Spinu V, Takahashi K, Vaughan D, Wilke C, Woo K and Yutani H** (2019) Welcome to the Tidyverse. *Journal of Open Source Software* **4**, 1686. doi:10.21105/joss.01686
- Woolhouse MEJ and Gowtage-Sequeria S** (2005) Host range and emerging and reemerging pathogens. *Emerging Infectious Diseases* **11**, 1842–1847. doi:10.3201/eid1112.050997
- Žiegytė R, Bernotienė R and Palinauskas V** (2022) *Culicoides segnis* and *Culicoides pictipennis* biting midges (Diptera, Ceratopogonidae), new reported vectors of *Haemoproteus* parasites. *Microorganisms* **10**, 898. doi:10.3390/microorganisms10050898