















- Karmakar, A. Lombardo, M. Bruna, L. N. Pfeiffer, K. W. West, A. C. Ferrari, M. Polini, and V. Pellegrini, Anomalous low-temperature Coulomb drag in graphene-GaAs heterostructures, *Nat. Commun.* **5**, 5824 (2014).
- [2] For a recent reference, see S. T. Chui, *Physics of the Electron Solid* (International Press, Boston, 1994); S. T. Chui and K. Esfarjani, A mechanism for quantum melting in two dimensions, *Europhys. Lett.* **14**, 361 (1991).
- [3] S. T. Chui and B. Tanatar, *Phys. Rev. Lett.* **74**, 458 (1995).
- [4] Y. Zhou, J. Sung, E. Brutschea, I. Esterlis, Y. Wang, G. Scuri, R. J. Gelly, H. Heo, T. Taniguchi, K. Watanabe *et al.*, Bilayer Wigner crystals in a transition metal dichalcogenide heterostructure, *Nature (London)* **595**, 48 (2021).
- [5] M. Huang, Z. Wu, N. Wang, and S. T. Chui, Quantum solid phase and Coulomb drag in two-dimensional electron-electron bilayers of MoS<sub>2</sub>, *Adv. Electron. Mater.* **9**, 2201105 (2023); S. T. Chui, M. Huang, Z. Wu, and N. Wang, Quantum structural transformation of correlation induced electron solids in two-dimensional MoS<sub>2</sub> double layers, *Phys. Rev. B* **108**, 165148 (2023).
- [6] E. P. Wigner, Effects of the electron interaction on the energy levels of electrons in metals, *Trans. Faraday Soc.* **34**, 678 (1938); See Eq. (11); See also E. Wigner, On the interaction of electrons in metals, *Phys. Rev.* **46**, 1002 (1934).
- [7] J. M. Kosterlitz and D. Thouless, Ordering, metastability and phase transitions in two-dimensional systems, *J. Phys. C* **6**, 1181 (1973); J. M. Kosterlitz, The critical properties of the two-dimensional xy model, *ibid.* **7**, 1046 (1974).
- [8] N. D. Mermin, The topological theory of defects in ordered media, *Rev. Mod. Phys.* **51**, 591 (1979); V. Poénaru and G. Toulouse, The crossing of defects in ordered media and the topology of 3-manifolds, *J. Phys. France* **38**, 887 (1977).
- [9] S. T. Chui and K. Esfarjani, Finite-temperature two-dimensional Wigner transition, *Phys. Rev. B* **44**, 11498 (1991).
- [10] S. Narasimhan and T.-L. Ho, Wigner-crystal phases in bilayer quantum Hall systems, *Phys. Rev. B* **52**, 12291 (1995).
- [11] F. Rapisarda and G. Senatore, Diffusion Monte Carlo study of electrons in two-dimensional layers, *Aust. J. Phys.* **49**, 161 (1996); F. Rapisarda and S. Conti, G. Senatore, Novel electron GAS systems, *Int. J. Mod. Phys. B* **13**, 479 (1999);
- [12] G. Goldoni, V. Schweigert, and F. M. Peeters, Stability and dynamical properties of a double-layer Wigner crystal in two dimensions, *Surf. Sci.* **361-362**, 163 (1996); G. Goldoni and F. M. Peeters, Stability, dynamical properties, and melting of a classical bilayer Wigner crystal, *Phys. Rev. B* **53**, 4591 (1996).
- [13] A. D. Migone and Z. R. Li, Melting transition of submonolayer Ar adsorbed on graphite, *Phys. Rev. Lett.* **53**, 810 (1984).
- [14] A. D. Novaco and J. P. McTague, *Phys. Rev. Lett.* **38**, 1286 (1977).
- [15] J. D. Jackson, *Classical Electrodynamics* (Wiley, New York, 1998).
- [16] A. Zangwill, *Modern Electrodynamics* (Cambridge University Press, New York, 2013), Section 8.3.
- [17] E. J. Mele, Screening of a point charge by an anisotropic medium: Anamorphoses in the method of images, *Am. J. Phys.* **69**, 557 (2001).
- [18] L. V. Keldysh, Coulomb interaction in thin semiconductor and semimetal films, *Sov. Phys. JETP Lett.* **29**, 658 (1979).
- [19] S. T. Chui, N. Wang, and C. Y. Wan, Quantum exciton solid in bilayer two-dimensional electron-hole systems, *Phys. Rev. B* **102**, 125420 (2020).
- [20] S. T. Chui, N. Wang, and B. Tanatar, Metal-insulator transitions in bilayer electron-hole systems in transition metal dichalcogenides, *Phys. Rev. B* **104**, 195432 (2021).
- [21] For some examples, See L. Bonsall and A. Maraduddin, Some static and dynamical properties of a two-dimensional Wigner crystal, *Phys. Rev. B* **15**, 1959 (1977); See the first terms on the right hand sides of Equations 2-15 to 2.21.
- [22] S. T. Chui and K. Esfarjani, Finite-temperature transport of a pinned 2D electron lattice, *Phys. Rev. Lett.* **66**, 652 (1991).
- [23] The top gate voltage changed by 1 V as the temperature is changed. This corresponded to a density changed of 3%. We have neglected this density change in our estimate.
- [24] S. T. Chui, Grain boundary theory of melting, *Phys. Rev. B* **28**, 178 (1983).
- [25] V. Pudalov and S. T. Chui, Character of the phase transition at the 2D electron liquid to solid boundary, *Phys. Rev. B* **49**, 14062 (1994).
- [26] I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series and Products* (Academic Press, New York, 2000), Eq. (6.611),